

AN OVERVIEW OF GENERALIZED BASIC LOGIC ALGEBRAS

PETER JIPSEN

Chapman University, California, USA

EXTENDED ABSTRACT

Classical propositional logic is one of the earliest formal systems of logic, with its origins in the work of Boole and De Morgan. The algebraic semantics of this logic is given by Boolean algebras. Both, the logic and the algebraic semantics have been generalized in many directions over the last 150 years. In this talk we primarily take the algebraic point of view, but we will also use the powerful framework of algebraic logic to clarify the close relationship between algebra and logic.

In various applications (such as fuzzy logic) the properties of Boolean conjunction are too stringent, hence a new binary connective \cdot , usually called *fusion*, is introduced. In Boolean algebra the relationship between conjunction and implication is given by the *residuation equivalences*

$$x \wedge y \leq z \iff x \leq y \rightarrow z \iff y \leq x \rightarrow z.$$

These equivalences imply many of the properties of \wedge and \rightarrow (such as commutativity of \wedge , distributivity of \wedge over \vee , left-distributivity of \rightarrow over \vee and right-distributivity of \rightarrow over \wedge) so one wishes to retain some aspects of these equivalences, but with conjunction replaced by fusion. To avoid imposing commutativity on the fusion operation one has to introduce two implications:

$$x \cdot y \leq z \iff x \leq y \rightarrow z \iff y \leq x \rightsquigarrow z.$$

Since the residuals $\rightarrow, \rightsquigarrow$ are generalized division operations, it is convenient to use the following alternative notation:

$$x \cdot y \leq z \iff x \leq z/y \iff y \leq x \setminus z.$$

Thus \setminus is another name for \rightsquigarrow , and $x \rightarrow y = y/x$. One advantage is that this notation allows the formulation of a simple *mirror image principle*: Any statement about residuated structures has an equivalent mirror image obtained by reading terms backwards (i.e. replacing $x \cdot y$ by $y \cdot x$ and interchanging x/y with $y \setminus x$), hence it suffices to state results in only one form.

In the most general setting a *residuated poset* $\langle P, \cdot, \setminus, /, \leq \rangle$ is a partially ordered set $\langle P, \leq \rangle$ with three binary operations that satisfy the above equivalences. In many applications to logic there exists additional structure, such as a constant 1 to denote *true*, and suprema and infima for finite subsets of P . Since fusion is usually assumed to be at least associative, we settle on a common foundation of Dilworth's *residuated lattices* [Di39], i.e., algebras of the form $\langle L, \vee, \wedge, \cdot, 1, \setminus, / \rangle$ such that

- $\langle L, \vee, \wedge \rangle$ is a lattice (i.e. \vee, \wedge are commutative, associative and mutually absorbentive),
- $\langle L, \cdot, 1 \rangle$ is a monoid (i.e. \cdot is associative, with identity element 1),
- $xy \leq z \iff x \leq z/y$ and $xy \leq z \iff y \leq x \setminus z$ for all $x, y, z \in L$.

Since the first two properties are defined by identities, it is important to note that this can also be done for the third: the equivalences are captured by $x(x \setminus z \wedge y) \leq z$, $y \leq x \setminus (xy \vee z)$ and their mirror images. Note that in the absence of parenthesis, we assume that \cdot is performed first, followed by $\setminus, /$ and finally \vee, \wedge . Moreover we use the notation $s \leq t$ as abbreviation for the equivalent identity $s = s \wedge t$. Hence residuated lattices form a *variety* (also called *equational class*), denoted by RL. A general introduction to varieties of universal algebras can be found in [BS81].

Note that in particular we do **not** assume that L is bounded or that 1 is the top element or that \cdot is commutative. These additional assumptions are handled by expanding the language with an additional constant 0, and/or adding further identities.

A residuated lattice with a constant 0 (which can denote any element) is called a *full Lambek algebra* or FL-algebra for short, and the variety of all such algebras is denoted by FL. A good introduction to FL and its associated logic can be found in [On03].

Much of the research on generalizations of BL-algebras can be viewed as investigations of certain subvarieties of FL and RL. We list here some of the important ones, together with their various names.

Some subvarieties of FL.

- FL_w -algebras [On03]: FL-algebras that satisfy $0 \leq x$ and $x \leq 1$.
- FL_e -algebras [On03]: FL-algebras that satisfy $x \cdot y = y \cdot x$ (or equivalently $x \setminus y = y/x$). In this case one usually writes $x \rightarrow y$ instead of $x \setminus y$ or y/x .
- DFL = *distributive FL-algebras*: FL-algebras that satisfy $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$.
- RFL = *representable FL-algebras*: FL-algebras that are subdirect products of linearly ordered FL-algebras, or equivalently satisfy the identity $1 \leq u \setminus ((x \vee y) \setminus x)u \vee v((x \vee y) \setminus y)/v$.
- psMTL = *pseudo monoidal t-norm algebras, or weak-pseudo-BL algebras* [FGI01]: FL_w -algebras that satisfy prelinearity ($x \setminus y \vee y \setminus x = 1$ and $x/y \vee y/x = 1$).
- FL_{ew} -algebras [KO01]: algebras that are both FL_e -algebras and FL_w -algebras.
- MTL = *monoidal t-norm algebras* [EG01]: FL_{ew} -algebras that satisfy prelinearity.
- psBL = *pseudo BL-algebras* [FGI01], [DGI02]: FL_w -algebras that satisfy divisibility ($x \wedge y = x(x \setminus y) = (y/x)x$).
- BL = *basic logic algebras* [Ha98]: MTL-algebras that satisfy divisibility, or equivalently, commutative prelinear pseudo BL-algebras.
- HA = *Heyting algebras* [BD74]: FL-algebras that satisfy $x \wedge y = xy$, or equivalently FL_w -algebras that are idempotent ($xx = x$).
- psMV = *pseudo MV-algebras* [GI01]: pseudo BL-algebras that satisfy $x \vee y = x/(y \setminus x) = (x/y) \setminus x$.
- MV = *multi-valued logic algebras, or Lukasiewicz algebras* [CDM00]: BL-algebras that satisfy $\neg \neg x = x$ or equivalently, commutative pseudo MV-algebras.
- GA = *Gödel logic algebras, or linear Heyting algebras* [Ha98]: BL-algebras that satisfy $x \cdot x = x$ or equivalently, Heyting algebras that satisfy prelinearity.

- **GMV** = *generalized MV-algebras* [JT02] [Ga03]: Residuated lattices that satisfy $x \vee y = x / ((x \vee y) \backslash x) = (x / (x \vee y)) \backslash x$.
- **Fleas** [Ha]: Integral residuated lattices that satisfy prelinearity ($x \backslash y \vee y \backslash x = 1$ and $x / y \vee y / x = 1$).
- **GBH** = **IGBL** = *generalized basic hoops, pseudo hoops*: Residuated lattices that satisfy $x \wedge y = x(x \backslash y) = (y / x)x$. It follows that they are integral.
- **BH** = *basic hoops* [AFM]: Commutative representable generalized basic hoops, i. e., commutative representable residuated lattices that satisfy divisibility ($x \wedge y = x(x \backslash y)$).
- **WH** = *Wajsberg hoops*: Commutative integral generalized MV-algebras.
- **CanRL** = *cancellative residuated lattices* [BCGJT]: Residuated lattices that satisfy $x = (x / y)y = y(y \backslash x)$. These identities are equivalent to cancellativity of fusion.
- **LG** = *lattice-ordered groups or ℓ -groups* [AF88] [GH89]: Residuated lattices that satisfy $1 = x(x \backslash 1)$. **NLG** = *normal-valued ℓ -groups*, defined by $(x \wedge 1)^2(y \wedge 1)^2 \leq (y \wedge 1)(x \wedge 1)$. **RLG** = *representable ℓ -groups*, defined by $1 \leq (e \backslash x)yx \vee 1 \backslash y$. **CLG** = *commutative ℓ -groups*.
- **LG⁻** = *negative cones of lattice-ordered groups* [JT02]: Cancellative integral generalized BL-algebras. **NLG⁻** = *negative cones of normal-valued ℓ -groups*, defined by $x^2y^2 \leq yx$ relative to **LG⁻**. **RLG⁻** = *negative cones of representable ℓ -groups*, defined as cancellative integral representable generalized BL-algebras. **CLG⁻** = *negative cones of commutative ℓ -groups*, defined as cancellative basic hoops.
- **Br** = *Brouwerian algebras*: Residuated lattices that satisfy $x \wedge y = xy$. **RBr** = *representable Brouwerian algebras*: Brouwerian algebras that satisfy prelinearity, or equivalently, basic hoops that are idempotent ($xx = x$).
- **GBA** = *generalized Boolean algebras*: Brouwerian algebras that satisfy $x \vee y = (x \rightarrow y) \rightarrow y$, or equivalently, Wajsberg hoops that are idempotent ($xx = x$). **GBA_n** = *generalized Boolean algebras of degree n*, defined as subdirect products of the linearly ordered $n+2$ -element Brouwerian algebra.

Many further varieties can be obtained from these by combining some of the identities mentioned above. For example the prefixes **C**, **D**, **I**, are used to denote the commutative, distributive and integral identities respectively. The relationships between various subvarieties of **RL** and **FL** are shown in Figures 1 and 2. Note that joins in these figures do not in general agree with joins in the lattice of subvarieties.

As indicated in Table 1, there is a close correspondence between certain subvarieties of **FL** and **RL**. In logic it is quite usual to have a constant 0 in the language to denote *falsity*. From an algebraic perspective it is in some ways natural to consider the slightly less expressive signature without 0 since, for example, the variety of ℓ -groups is not a subvariety of **FL**.

The notion of triangular norm has been studied extensively in the theory of probabilistic measures. Since they give rise to generating algebras for several subvarieties, we recall the definition here. A *pseudo-t-norm* is an order-preserving monoid operation on the unit interval $[0, 1]$, with 1 as the identity. A *t-norm* is a commutative pseudo-t-norm. It is said to be continuous if it is a continuous function from $[0, 1]^2$ to $[0, 1]$ in the standard topology of the unit interval. Table 2 summarizes which varieties are generated by algebras based on specific t-norms.

FL	RL	Defining identities
FL_e	CRL	$xy = yx$
DFL	DRL	$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$
RFL	RL^C	$1 \leq u \setminus ((x \vee y) \setminus x)u \vee v((x \vee y) \setminus y)/v$
		Below add $0 \leq x$ for subvarieties of FL
FL_w	IRL	$x \leq 1$
FL_{ew}	CIRL	$xy = yx, x \leq 1$
psMTL	Fleas	$x \setminus y \vee y \setminus x = 1 = x/y \vee y/x$
MTL	$CIRL^C$	$xy = yx, (x \rightarrow y) \vee (y \rightarrow x) = 1$
	GBL	$x \wedge y = x(x \setminus (x \wedge y)) = ((x \wedge y)/x)x$
psBL	IGBL	$x \wedge y = x(x \setminus y) = (y/x)x$
BL	BH	$xy = yx, x \wedge y = x(x \rightarrow y), (x \rightarrow y) \vee (y \rightarrow x) = 1$
	GMV	$x \vee y = x / ((x \vee y) \setminus x) = (x / (x \vee y)) \setminus x$
psMV	IGMV	$x \vee y = x / (y \setminus x) = (x/y) \setminus x$
MV	WH	$xy = yx, x \vee y = (x \rightarrow y) \rightarrow y$
HA	Br	$x \wedge y = xy$
GA	RBr	$x \wedge y = xy, (x \rightarrow y) \vee (y \rightarrow x) = 1$
Π		BL and $\neg\neg x \leq (x \rightarrow xy) \rightarrow y(\neg\neg y)$
BA	GBA	$x \wedge y = xy, x \vee y = (x \rightarrow y) \rightarrow y$

TABLE 1. Definition and correspondence of subvarieties of FL and RL

FL	RL	Generated by
MTL	$CIRL^C$	all residuated t-norms [JM02]
BL	BH	all continuous t-norms [Ha98] [CEGT00]
MV	WH	Lukasiewicz $xy = \max\{0, x + y - 1\}$ [Ch59]
GA	RBr	Gödel $xy = \min\{x, y\}$ [Ha98]
Π		Product $xy = \text{multiplication on } [0, 1]$ [Ha98]

TABLE 2. Some varieties generated by t-norms

The subvarieties of RL are obtained if we do not specify 0 as a constant of the algebra.

The relationship between subvarieties of FL and RL is illuminated by the following reduct and expansion functors: $R_0 : FL \rightarrow RL$ the forgetful functor that removes 0 from the signature, and $E_0 : RL \rightarrow FL$ given by $E_0(A) = 0 \oplus A$, the ordinal sum of $\{0\}$ with A on top, and $0x = x0 = 0$. Given a subvariety \mathcal{V} of FL, we obtain a subvariety of RL by defining $R_0\mathcal{V}$ to be the variety generated by $\{R_0A \mid A \in \mathcal{V}\}$. Similarly, a subvariety \mathcal{W} of RL is mapped to a subvariety of FL by defining $E_0\mathcal{W}$ to be the variety generated by $\{E_0B \mid B \in \mathcal{W}\}$. Both maps preserve inclusions of varieties.

Any subvariety of RL or FL has a corresponding logic that is sound and complete with respect to the algebraic semantics of the subvariety. Thus many logical questions have algebraic counterparts and vice versa.

In the algebraic setting there are some semantic methods that can shed light on the logical side. For example one can study the subdirectly irreducible members of the variety, or ask about the finite embedding property [BvA02] [BvA], equationally definable principal congruences (EDPC) [BP82], and the existence of a discriminator term [Jo95].

We will see how the congruences in residuated lattices are ideal determined, correspond to convex normal subalgebras, and how this characterization is used to give equational bases for some subvarieties [GU84] [JT02] [vA02]. EDPC is related to the deduction theorem in logic, and discriminator varieties are equivalent to expansions with Baaz's projection Δ [Ba96].

In the logical setting, there are syntactic methods such as cut-free Gentzen systems that have interesting consequences on the algebraic side, for example decidability [OK85] [OT99] and joint embedding properties.

Finally, we will consider the categorical equivalence between lattice ordered groups and cancellative integral generalized BL-algebras [JT02] [BCGJT] [vA], as a special case of the very general Morita equivalence in universal algebra [McK96]. Further we will mention a recent generalization of Mundici [Mu86] and Dvurečenskij's [Dv02] categorical equivalences regarding (pseudo) MV-algebras to an equivalence between lattice ordered groups expanded with a nucleus and generalized MV-algebras. The latter result is due to N. Galatos [Ga03] where he used it to prove the decidability of the equational theory of generalized MV-algebras. A recent implementation, using the decision procedure for lattice ordered groups, due to Holland and McCleary [HM79], can be found at [www.chapman.edu/~jipsen/].

REFERENCES

- [AFM] P. Agliano, I. M. A. Ferreira and F. Montagna, *Basic hoops: An algebraic study of continuous t-norms*, preprint.
- [AF88] M. Anderson and T. Feil, *Lattice-Ordered Groups: an introduction*, D. Reidel Publishing Company, 1988.
- [Ba96] M. Baaz, *Infinite-valued Gödel logics with 0-1-projections and relativizations*, in: Gödel 1996, Brno, Springer, Berlin (1996), 23–33.
- [BCGJT] P. Bahls, J. Cole, N. Galatos, P. Jipsen, C. Tsinakis, *Cancellative residuated lattices*, Algebra Universalis, to appear.
- [BD74] R. Balbes and P. Dwinger, *Distributive Lattices*, University of Missouri Press, Columbia, Mo., 1974.
- [BJO] F. Belardinelli, P. Jipsen and H. Ono, *Algebraic aspects of cut elimination*, preprint.
- [Bi67] G. Birkhoff, *Lattice Theory*, (3rd ed), Colloquium Publications 25, Amer. Math. Soc., 1967.
- [BP82] W. J. Blok and D. Pigozzi, *On the structure of varieties with equationally definable principal congruences I*, Algebra Universalis 15, 195–227.
- [BvA02] W. J. Blok and C. J. van Alten, *The finite embeddability property for residuated lattices, pocrim's and BCK-algebras*, Algebra Universalis 48, (2002), 253–271.
- [BvA] W. J. Blok and C. J. van Alten, *The finite embeddability property for partially ordered biresiduated integral groupoids*, preprint.
- [BT] K. Blount and C. Tsinakis, *The structure of residuated lattices*, Intl. Journal of Algebra and Computation, to appear.
- [BS81] S. Burris and H. P. Sankappanavar, *A Course in Universal Algebra*, Springer Verlag, 1981, online at <http://www.thoralf.uwaterloo.ca/>

- [Ch59] C. C. Chang, *A new proof of the completeness of the Łukasiewicz axioms*, Trans. Amer. Math. Soc. 93 1959 74–80.
- [CDM00] R. Cignoli, I. D’Ottaviano, D. Mundici, *Algebraic foundations of many-valued reasoning*, Trends in Logic—Studia Logica Library, 7. Kluwer Academic Publishers, Dordrecht, 2000.
- [CEGT00] R. Cignoli, F. Esteva, L. Godo and A. Torrens, *Basic fuzzy logic is the logic of continuous t -norms and their residua*, Soft Computing 4, (2000), 106–112.
- [Ci01] P. Cintula, *About axiomatic systems of product fuzzy logic*, Soft Computing 5, (2001), no. 3, 243–244.
- [DGI02] A. Di Nola, G. Georgescu and A. Iorgulescu, *Pseudo-BL algebras I*, Mult. Val. Log. 8 (2002), no. 5-6, 673–714.
- [Di39] R. P. Dilworth, *Non-commutative residuated lattices*, Trans. Amer. Math. Soc. 46, (1939), 426–444.
- [Dv02] A. Dvurečenskij, *Pseudo MV-algebras are intervals in l -groups*, J. Aust. Math. Soc. 72 (2002), no. 3, 427–445.
- [EG01] F. Esteva and L. Godo, *Monoidal t -norm based logic: towards a logic for left-continuous t -norms*, Fuzzy Sets and Systems 124 (2001), no. 3, 271–288.
- [Fe92] I. M. A. Ferreira, *On varieties and quasivarieties of hoops and their reducts*, Ph. D. thesis, University of Illinois at Chicago, USA, 1992.
- [FGI01] P. Flondor, G. Georgescu and A. Iorgulescu, *Pseudo- t -norms and pseudo-BL-algebras*, Soft Computing 5 (2001), 355–371.
- [Ga03] N. Galatos, *Varieties of residuated lattices*, Ph. D. thesis, Vanderbilt University, Nashville, USA, 2003.
- [GH89] A. M. W. Glass and W. C. Holland (editors), *Lattice-Ordered Groups*, Kluwer Academic Publishers, 1989, 278–307.
- [GI01] G. Georgescu and A. Iorgulescu, *Pseudo-MV algebras*, Mult. Val. Logic 6 (2001), 95–135.
- [GU84] H.P. Gumm and A. Ursini, *Ideals in universal algebras*, Algebra Universalis 19 (1984), no. 1, 45–54.
- [Ha98] P. Hájek, *Metamathematics of Fuzzy Logic*, volume 4 of *Trends in Logic*, Kluwer Academic Publishers, 1998.
- [Ha02] P. Hájek, *Observations on the monoidal t -norm logic*, Fuzzy Sets and Systems 132 (2002), no. 1, 107–112.
- [Ha] P. Hájek, *Fleas and fuzzy logic*, preprint.
- [HRT02] J. Hart, L. Rafter and C. Tsinakis, *The structure of commutative residuated lattices*, International Journal of Algebra and Computation, 12 (2002), no. 4, 509–524.
- [HM79] W. C. Holland and S. H. McCleary, *Solvability of the word problem in free lattice-ordered groups*, Houston J. Math. 5 (1979), no. 1, 99–105.
- [JM02] S. Jenei and F. Montagna, *A proof of standard completeness for Esteva and Godo’s logic MTL*, Studia Logica 70 (2002), no. 2, 183–192.
- [JT02] P. Jipsen and C. Tsinakis, *A Survey of Residuated Lattices*, in “Ordered Algebraic Structures” (J. Martinez, editor), Kluwer Academic Publishers, Dordrecht, 2002, 19–56.
- [Jo95] B. Jónsson, *Congruence distributive varieties*, Math. Japonica 42 (1995), no. 2, 353–401.
- [KO01] T. Kowalski and H. Ono, *Residuated lattices: An algebraic glimpse at logics without contraction*, monograph, March 2001, 1–67.
- [KO00] T. Kowalski and H. Ono, *Splittings in the variety of residuated lattices*, Algebra Universalis 44 (2000), no. 3-4, 283–298.
- [McK96] R. McKenzie, *An algebraic version of categorical equivalence for varieties and more general algebraic categories*, Logic and Algebra (ed. A. Ursini, P. Aglianò), Marcel Dekker, 1996, 211–244.
- [Mu86] D. Mundici, *Interpretation of $AF C^*$ -algebras in Łukasiewicz sentential calculus*, J. Funct. Anal. 65 (1986), no. 1, 15–63.
- [On03] H. Ono, *Substructural Logics and Residuated Lattices — an Introduction*, Studia Logica, to appear.
- [OK85] H. Ono and M. Komori, *Logics without the contraction rule*, Journal of Symbolic Logic, 50 (1985) 169–201.

- [OT99] M. Okada and K. Terui, *The finite model property for various fragments of intuitionistic linear logic*, Journal of Symbolic Logic, **64**(2) (1999) 790–802.
- [vA02] C. J. van Alten, *Representable biresiduated lattices*, Journal of Algebra 247 (2002), no. 2, 672–691.
- [vA] C. J. van Alten, *The termwise equivalence of the varieties of ℓ -group cones and generalized cancellative hoops*, preprint.
- [Wa40] M. Ward, *Residuated distributive lattices*, Duke Math. J. 6, (1940). 641–651.
- [WD38] M. Ward and R.P. Dilworth, *Residuated Lattices*, Proceedings of the National Academy of Sciences 24 (1938), 162–164.
- [WD39] M. Ward and R.P. Dilworth, *Residuated Lattices*, Transactions of the AMS 45 (1939), 335–354.